Ruptured Inferior Mesenteric Artery Aneurysm: Case Report

Herzallah AM; Consultant vascular surgeon; Matarya Teaching Hospital; Cairo; Egypt.

Abstract (300 word limit)

Among visceral artery aneurysms, inferior mesenteric artery aneurysm is the rarest. While most cases are of degenerative aetiology, a small fraction of these cases is caused by autoimmune diseases. We report a case of spontaneous retroperitoneal hemorrhage from a ruptured IMA aneurysm in a 30 years old patient with SLE, treated by ligation.

.

Recent Publications (minimum 5)

1. **Stanley JC, Wakefield TW, Graham LM, WhitehouseWMJr, Zelenock GB, Lindenauer SM.:** Clinical importance and management of splanchnic artery aneurysms. J Vasc Surg 1986;3:836-40

2. **Shanley, CJ., Shah, NL,. and Messina, LM.:** Uncommon Splanchnic Artery Aneurysms: Pancreaticoduodenal, Gastroduodenal, Superior Mesenteric, Inferior Mesenteric, and Colic. Annal Vasc Surg 1996; 10(5): 506-15

3. **Parvin, SD., Earnshaw, JJ., & Joint Vascular Research Group:** Rare vascular disorders: A practical guide for the vascular specialist 2005; 196

4. **Ko. SF., Hsien. MJ., NG. SH. et al.: Superior mesenteric artery aneurysm in systemic lupus erythematosus. Clinical Imaging 1997; 21(1): 13-16**

5. **Edogawa, S., Shibuya, T., Kurose, K., Sasaki, K., & Tomita, H.:** Inferior mesenteric artery aneurysm: case report and literature review. Annals of vascular diseases 2013; 6(1), 98-101

6. **Tovar, JR., Molina, EM., Morales, V., Sanjuanbenito, A. and Lobo, E:** Evolution of the therapeutic approach of visceral artery aneurysms. Scandinavian Journal of Surgery 96: 308–313, 2007



Biography (150 word limit)

. **Herzallah AM; Consultant vascular surgeon; Matarya Teaching Hospital; Cairo; Egypt.**

Email:assemherz010@hotmail.com

Notes/Comments: