Abstract Title:

Three-dimensional culture system by magnetic levitation for the study of hematopoietic stem cell microenvironment

*[1]Mejía-Cruz CC, [1]Barreto E, [1]Pardo-Pérez MA, [2]Leal-García E, [2]Pérez-Núñez R, [3]Jaramillo LF, [3]Rodríguez-Sarmiento JL, [1]Barreto A, [1]**Rodríguez-Pardo VM.***

[1] Grupo de Inmunobiología y Biología Celular. Departamento de Microbiología. Facultad de Ciencias. Pontificia Universidad Javeriana, Carrera 7a # 43-82, Edificio Félix Restrepo (Edificio 50). Bogotá D.C., Colombia (Sur America).
[2] Departamento de Ortopedia y Traumatología. Facultad de Medicina. Pontificia Universidad Javeriana y Hospital Universitario San Ignacio, Carrera 7a # 40-62. Bogotá D.C., Colombia (Sur America).
[3] Departamento de Patología. Facultad de Medicina. Pontificia Universidad Javeriana y Hospital Universitario San Ignacio, Carrera 7a # 40-62. Bogotá D.C., Colombia (Sur America).

Exhibitor author: Viviana Marcela Rodríguez-Pardo
e-mail: vivianar@javeriana.edu.co or vivianarodrigu@gmail.com
Grupo de Inmunobiología y Biología Celular. Departamento de Microbiología. Facultad de Ciencias. Pontificia Universidad Javeriana, Carrera 7a # 43-82, Edificio Félix Restrepo (Edificio 50). Bogotá D.C., Colombia (Sur America). PBX (571-3208320 – 4072)

Abstract (not more than 300 words)

Background. Hematopoietic stem cells (HSC) proliferation, differentiation, self-renewal and survival are regulated by specific characteristics of their microenvironment. Bone marrow (BM), HSC's niche, is composed by different cell populations that by their signaling and interactions regulate HSC. Attempts to recreate this complex microenvironment have been devised in various studies using diverse co-culture models with different types of cell populations. The objective of this study is to propose a 3D magnetic levitation culture, free of any exogenous structures and substances, in which HSC are co-cultured with mesenchymal stem cells (MSC) and endothelial cells (EC), where a multicellular sphere is formed, providing an organoid model to be analyzed. **Methods.** For the conformation of the multicellular spheres, MSC were isolated from human BM, HSC from human umbilical cord blood and EC line Lonza CC.2811 was used. System standardization was accomplished by size and shape spheres evaluation; percentage of cell aggregation; viability assay and HSC CFU-assay; followed by immunohistochemical and immunofluorescent analysis of histological sections. **Results.** Nanoparticles density was stablished at 1uL /10.000 cells. The optimum ratio for the culture of the three cell populations was 1 MSC: 2 HSC: 2 EC. Multicellular spheres were completely formed at day 10 on culture, accomplishing a sphericity >0.8 and an aggregation percentage of 70-90% on day 5. The HSC isolated form the multicellular spheres preserve their multipotent function. **Discussion**. The multicellular spheres obtained with this methodology are very suitable in evaluating proliferation, differentiation, clonogenic potential and other aspects of HSC when in contact with other BM cell populations, thus allowing for further studies to evaluate how HSC and its niche respond to different treatments, drugs and stress caused by

infections, myelosuppression, neoplasms and aging, among other factors, so that future therapies for hematological diseases can be developed.

Key words: Magnetic levitation system, Hematopoietic stem cells, Bone marrow, Microenvironment, Mesenchymal stem cells

Professional Biography (100-150 words)

Viviana Marcela Rodríguez-Pardo (exhibitor author). I am the leader of the research line of "Biology of stem cells," in the Immunobiology and Cellular Biology group at the Faculty of Sciences of the Pontificia Universidad Javeriana (www.javeriana.edu.co) in the city of Bogota (Colombia, South America) during 12 years. During my activity as a researcher, I directed different research projects that have allowed me to develop protocols for isolation, in vitro expansion and characterization of human hematopoietic stem cells (1, 2) and human mesenchymal stem cells (3, 4) to solve problems related to the modulation of stem cells by their microenvironment. In addition, I have participated with other researchers on projects related to the role of the stem cells in the tumor microenvironment (3, 5).

1. Rodriguez-Pardo VM, Vernot JP. Mesenchymal stem cells promote a primitive phenotype CD34+c-kit+ in human cord blood-derived hematopoietic stem cells during ex vivo expansion. Cell Mol Biol Lett. 2013;18(1):11-33.
2. Pardo-Pérez MJ, Mejía-Cruz C, Leal-García E, Pérez-Núñez R, Useche LF, Ulloa JC, Jaramillo LF, Rodríguez-Pardo VM. . Three-dimensional tissue culture system by the magnetic levitation method for the study of the human bone marrow microenvironment. Tissue Engineering. Dic 2017 - Submitted.
3. Rodriguez-Pardo VM, Aristizabal JA, Jaimes D, Quijano SM, de los Reyes I, Herrera MV, et al. Mesenchymal stem cells promote leukaemic cells aberrant phenotype from B-cell acute lymphoblastic leukaemia. Hematol Oncol Stem Cell Ther. 2013;6(3-4):89-100.
4. Rodríguez-Pardo VM FLM, Aristizabal JA, Vernot JP. Aislamiento y caracterización de células "stem" mesenquimales de médula ósea humana según criterios de la Sociedad Internacional de Terapia Celular Universitas Scientiarum. 2010;15(3):224-39.
5. Vernot JP, Bonilla X, Rodriguez-Pardo V, Vanegas NP. Phenotypic and Functional Alterations of Hematopoietic Stem and Progenitor Cells in an In Vitro Leukemia-Induced Microenvironment. Int J Mol Sci. 2017;18(2).



Email: vivianar@javeriana.edu.co
vivianarodrigu@gmail.com