Long Term Study on the Clinical Effectiveness of Wet Cupping in Patients suffering from Sciatic Pain

Corresponding Author : **Prof.Dr.Muhammad Amin Baig**
DSc (TM), PhD (USA), FRCP, MD (AM) Sri Lanka, MD(EH), PGD(TCM), FTJ, DHMS,
Chairman International Alternative Medical Council (IAMC) , Islamabad Pakistan
HoD & Professor of Unani Medicine & TCM, *at Ajmal Unani Medical College Rawalpindi*
Visiting Professor Open International University for Complementary Medicine , Sri-Lanka
(draminbaig1976@gmail.com)

Co Author : **Dr.Syed Muzzammil Masaud**(RPh) Pharm-D , M.phil ,
Director Academics International Alternative Medical Council (IAMC) Pakistan,
Assistant Professor Islamabad Institute of Health Sciences (IIHS) Islamabad ,
Pakistan.(dr.muzamil15@gmail.com)

Reviewed by: **Hakeem Kamran Latif Chughtai** Controller of Examinations, National Council for Tibb, Government of Pakistan

Abstract:

Background : Ancient old Technique of Wet Cupping commonly known as HIJAMA has proven its effectiveness in number of medical ailments including Sciatic Nerve Pain. Long Term study to evaluate the effectiveness of Wet Cupping after 2 years and 5 years for patients suffering from sciatic nerve pain.

Method :

250 patients were randomly selected having no previous physical injury, between age 18-80years.Wet cupping sessions were done on every alternate day, 3 sessions were done in which removal of subcutaneous blood with the help of Glass suction cups was done after making superficial incisions . One cup was generally applied Between shoulders at T1 and C7 vertebrae area, and Two Cups were applied on L3-L4 or L4-L5 and L5-S1 vertebrae . After these sessions the Decrease in Sciatic Pain and improvement in Quality of life of Patient was assessed by Visual analogue Score, Numeric Pain rating scale, and WHOQOL scale at before treatment , After Treatment, 1year after treatment, 2years after Treatment, and 5 years after treatment. (Data were analyzed by paired t test and ANOVA.)

Result:

Out of 250 patients about 225 (90%)of Patients have shown marked decrease in sciatic nerve pain. After 1year of treatment 175 (70%) of Patients were satisfied and doesn't need any further treatment of sciatic pain, after 2year of treatment more than 125 (50%) patients showed marked decrease in pain as compared to before treatment.5years after treatment 92(37%) of patients were enjoying good quality of life as shown by their WHOQOL scale.

Conclusion: Long term benefits of Wet Cupping in patients suffering from Sciatic nerve pain was observed , it is concluded that overall quality of life of patient is improved and pain score is significantly decreased after 2years and 5years of HIJAMA Treatment.

Keywords: Wet Cupping, HIJAMA, Sciatic Nerve Pain, WHOQOL scale.

References:

[1] Albinali Hajar, H.A. (2004) Arab Gulf Traditional Medicine-Blood Letting. Heart Views, 5, 178-183.

[2] Farhadi, K., Schwebel, D.C., Saeb, M., Choubsaz, M., Mohammadi, R. and Ahmadi, A. (2009) The Effectiveness of Wet-Cupping for Nonspecific Low Back Pain in Iran: A Randomized Controlled Trial. Complementary Therapies in Medicine, 17, 9-15. http://dx.doi.org/10.1016/j.ctim.2008.05.003

[3] Ahmadi, A., Schwebel, D.C. and Rezaei, M. (2008) The Efficacy of Wet-Cupping in the Treatment of Tension and Migraine Headache. American Journal of Chinese Medicine, 36, 37-44. http://dx.doi.org/10.1142/S0192415X08005564 M. Bilal, R. A. Khan 330

[4] Lauche, R., Cramer, H., Hohmann, C., et al. (2012) The Effect of Traditional Cupping on Pain and Mechanical Thresholds in Patients with Chronic Nonspecific Neck Pain: A Randomized Controlled Pilot Study. Evidence-Based Complementary and Alternative Medicine : eCAM, 429718. http://dx.doi.org/10.1155/2012/429718

[5] Valat, J.P., Genevay, S., Marty, M., Rozenberg, S. and Koes, B. (2010) Sciatica. Best Practice & Research. Clinical Rheumatology, 24, 241-252. http://dx.doi.org/10.1016/j.berh.2009.11.005

[6] Iyer, K.M. (2013) Back Pain and Slipped Intervertebral Disc. General Principles of Orthopedics and Trauma, Springer-Verlag, London, 243-272. http://dx.doi.org/10.1007/978-1-4471-4444-1_16

[7] Koes, B.W., Van Tulder, M.W. and Peul, W.C. (2007) Diagnosis and Treatment of Sciatica. BMJ, 334, 1313-1317. http://dx.doi.org/10.1136/bmj.39223.428495.BE

[8] Miller, T.A., White, K.P. and Ross, D.C. (2012) The Diagnosis and Management of Piriformis Syndrome: Myths and Facts. Canadian Journal of Neurological Sciences, 39, 577-583. http://dx.doi.org/10.1017/S0317167100015298

[9] Casey, E. (2011) Natural History of Radiculopathy. Physical Medicine and Rehabilitation Clinics of North America, 22, 1-5. http://dx.doi.org/10.1016/j.pmr.2010.10.001

[10] Kirschner, J.S., Foye, P.M. and Cole, J.L. (2009) Piriformis Syndrome, Diagnosis and Treatment. Muscle Nerve, 40, 10-18. http://dx.doi.org/10.1002/mus.21318

[11] Bruggeman, A.J. and Decker, R.C. (2011) Surgical Treatment and Outcomes of Lumbar Radiculopathy. Physical Medicine and Rehabilitation Clinics of North America, 22, 161-177. http://dx.doi.org/10.1016/j.pmr.2010.10.002

[12] El Sayed, S.M., Mahmoud, H.S. and Nabo, M.M.H. (2013) Medical and Scientific Bases of Wet Cupping Therapy (AlHijamah): In Light of Modern Medicine and Prophetic Medicine. Alternative & Integrative Medicine, 2, 5.

[13] Luijsterburg, P.A.J., Verhagen, A.P., Ostelo, R.W.J.G., Oston, A.G., Peul, W.C. and Koes, B.W. (2007) Effectiveness of Conservative Treatments for the Lumbosacral Radicular Syndrome: A Systematic Review. European Spine Journal, 16, 881-899