Elevated Circulating microRNAs and Vasoactive Amine Metabolites in Neurocardiogenic Syncope Patients

A PILOT SYUDY

Amira S. HABIB [a], Hazem M. WARDA[a, b], Ahmed WAGDI [c], Amany ELSHORBAGY [d], Ahmad R. BASSIOUNY [e]

a) Alhyatt Heart and Vascular Center, Alexandria, Egypt
b) Department of Cardiology, Faculty of Medicine, Tanta University Hospital, Tanta, Egypt
c) Institute of Pharmacology and Toxicology, University Medical Center Goettingen, Goettingen, Germany
d) Department of Physiology, Faculty of Medicine, University of Alexandria, Alexandria, Egypt
e) Department of Biochemistry ,Faculty of Science, University of Alexandria, Alexandria, Egypt

Author for correspondence: Amira S.Habib, Alhyatt Heart and Vascular Center, Alexandria, Egypt. **Email:** ahabib@alhyatt.com **Tel:** 002 0122 2396 706
Address: 90 Fawzy Moaz st., Alhyatt Heart and vascular center, Smouha, Alexandria, Egypt

ABSTRACT

OBJECTIVE

Syncope is a common clinical problem challenging both cardiologists and general practitioners. In our study, we selected a number of circulating micro-RNAs and vasoactive amine metabolites to be evaluated in attempts to introduce possible biomarkers for syncope.
To the best of our knowledge, this is the first study to assess the changes in miRNAs in syncope patients.

MATERIALS AND METHODS

Nineteen patients with history of syncope and nineteen sex and age matched healthy controls participated. A detailed medical history was recorded and a cardiovascular examination followed by head up tilt table testing (HUTT) were performed. Three blood samples were withdrawn, first one at baseline, second during syncopal attack and third one after 30 minutes from the end of the tilt test. The levels of three circulating microRNAs (miR-210, miR-1 and miR-34a) and three vasoactive amine metabolites (endothelin-1, copeptin and serotonin) were quantified.

RESULTS

In Group A, copeptin significantly increased during syncopal attack by 21.7 ± 0.45 pg/mL vs 4.3 ± 1.209 pg/mL in control subjects (Group B; P= 0.002). Similarly, endothelin-1 values significantly rose by an average of 28 ± 1.25 pg/mL in syncope patients versus 3.35 ± 0.75 pg/mL in healthy controls (P<0.001). While Serotonin (5-HT) levels were significantly greater during syncope relative to baseline in HUTT positive patients by 95.89 ± 3.7 pg/mL versus 9 ± 1.43 pg/ml (p<0.001) in control subjects. In summary, vasoactive amines increased with a 3-5 fold change in group A, but showed 1-2 fold increase in the control group.
In group A, miR-210 increased by mean of 0.6 ± 0.09 (p<0.001) during syncope (95% CI [0.4, 0.79]). While miR-34a values increased by mean of 0.89 ± 0.22 during syncope than baseline value (95% CI [0.42, 1.36]) with significant difference of P =0.001. Likewise, miR-1 levels was elevated by an average of 0.42 ± 0.07 (95% CI [0.26, 0.58]) with a p<0.001 significance. miRNA levels were 3±1 fold higher in the syncope patients (Group A) than in controls.

CONCLUSION

The selected miRNAs and vasoactive amines have a very promising diagnostic and therapeutic potential as biomarkers in diagnosing syncope.