5th International Conference on Advances in Skin, Wound Care and Tissue Science
October 15-16, 2018 Rome, Italy

The effect of cold atmospheric pressure plasma (CAP) on cell migratory behaviors and molecular markers of wound healing machinery

D. Shome[1], N. Hohmann[1], J. Schmidt[2], T. von Woedtke[1] and K. Masur[1]

[1]Leibniz Institute for Plasma Science and Technology e.V., Greifswald, Germany.

[2] Clinic Karlsburg, Dr. Guth GmbH & Co. KG, Karlsburg, Germany

Abstract:

Cold atmospheric pressure plasma (CAP) is a promising tool for biomedical and clinical application. Cold physical plasma are partially ionized gases that mediate biological response generating ROS and RNS species.[1-3]In this study, we used the medical device class 2a kINPen MED®; it is clinically approved atmospheric argon plasma jet. Typical active agents being generated include ions, electrons, and reactive oxygen and nitrogen species (ROS/RNS). Electric and magnetic fields, light (visible, in-frared, UV), and neutral particles are also being generated [4]. It is well known, that wound oxygenation is an important factor of wound healing and scavenging reactive species could impair wound healing.[5] In this study, we examined the extent of wound healing and the underlying cellular mechanism *in vitro* induced by CAP. Our group had also collected wound exudates before and after CAP treatment from 7 ambulant diabetic patients with chronic wounds from 'Competence Center Diabetes' in Karlsburg. These wound exudates were also analyzed.

Methods: For our studies, we incorporated dermal keratinocytes, fibroblasts and co culture. An indirect treatment where CAP treated media (RPMI) was added and a direct CAP treatment on the cells (different time points) were performed and the migration assay was monitored. Matrix metalloproteinase play a pivotal role in wound re epithelization. The amount of metalloproteinase and several cytokines (especially Interleukins) in the patient exudates were also monitored by ELISA.

Results and future direction: Short-term CAP treatment induces enhanced cell migration than longer treatment and compared to untreated control. Co culture studies show an improved cell migration upon CAP treatment compared to keratinocytes alone. Matrix metalloproteinases and Interleukins were also reduced in wound exudates after CAP treatment which indicates improved wound healing. It was also evident with an average of more than 80 % reduction in the wound size of the patients undergoing CAP treatment. Also, the signaling machinery of wound healing involving inflammatory (mediated by interleukins) and regenerative pathways (mediated by HIPPO signaling) are currently being checked by quantative PCR and western blot to identify an autocrine/paracrine signaling pathway induced by CAP.

References:

1. Jablonowski, H. and T. von Woedtke (2015). "Research on plasma medicine-relevant plasma–liquid interaction: What happened in the past five years?" Clinical Plasma Medicine 3(2): 42-52.

2. Reuter, S., et al. (2012). "From RONS to ROS: Tailoring Plasma Jet Treatment of Skin Cells." IEEE Transactions on Plasma Science 40(11): 2986-2993.

3 David, B. G. (2012). "The emerging role of reactive oxygen and nitrogen species in redox biology and some implications for plasma applications to medicine and biology." Journal of Physics D: Applied Physics 45(26): 263001.

4. Bekeschus, S., et al. (2016). The plasma jet kINPen – A powerful tool for wound healing. Clinical Plasma Medicine. 4: 19-28.

5. Schmidt, A., et al. (2017). "One Year Follow-Up Risk Assessment in SKH-1 Mice and Wounds Treated with an Argon Plasma Jet." International Journal of Molecular Sciences 18(4): 868.

Biography:

I, Miss Debarati Shome have graduated from St. Xavier's College, Kolkata, India with my major in Microbiology. Later, I obtained my post-graduation from University of Calcutta, India with my major in Biotechnology. Presently, I am working as a PhD student in Leibniz Institute for Plasma Science & Technology (INP), Greifswald in the Plasma Wound Healing (PWH) group. Here, I am focusing on the effect of Cold atmospheric pressure plasma in wound healing.

Keywords: Cold Atmospheric Pressure Plasma (CAP), Matrix Metalloproteinases (MMPs), kINPen MED®, Interleukins

Desired presentation: Oral/Poster

Topic: Track 13: Skin & Wound Healing

Corresponding Author Details: **1.** Debarati Shome
debarati.shome@inp-greifswald.de
Leibniz Institute for Plasma Science & Technology (INP), Greifswald
Felix Hausdorff Str. 2, 17489 Greifswald, Germany
Telephone: +49 3834 554 3864
Personal: +49 1744831021

2. Dr. Kai Masur
kai.masur@inp-greifswald.de
Group Leader, Plasma Wound Healing (PWH)
Leibniz Institute for Plasma Science & Technology (INP), Greifswald
Felix Haudorff Str 2, 17489 Greifswald
Telephone: +49 3834 554 3322