

EXOSOME NANOVESICLES FOR THE HOMING DELIVERY OF THERAPEUTICS INTO AUTOLOGOUS CANCER CELLS

Shuhua Bai Ph.D.

April 18, 2018


Associate Professor

**Department of Pharmaceutical Sciences
School of Pharmacy, Husson University**


Outline

- Introduction
 - Nanoparticle-enhanced chemotherapy
 - Exosome for targeted drug delivery
- Experiment
 - Exosome-homing evaluation
 - Exosome-homing delivery
 - Fluorescent marker and anticancer drugs
 - Small interfering RNA
 - Exosome characterization
 - Surface proteins and miRNA
- Conclusions
- Future studies

Targeted-Delivery of Chemotherapy





Traditional Chemotherapy



Targeted Chemotherapy


Nanoparticles as Delivery Carriers

- Free anticancer drugs loaded into carrier
- Protection
- Pharmacokinetics
 - Fusion
 - Endocytosis uptake




Nanoparticles-Enhanced Delivery

- Passive targeting
 - Enhanced permeability and retention (EPR)
- Limitation of nanoparticle-delivered anticancer drugs
 - Active targeting
 - Immunogenicity





Extracellular Vesicles

➤ Garbage bins to novel carriers




**2013 Nobel Prize
Physiology or Medicine**



Graça Raposo, and Willem Stoorvogel , J Cell Biology 2013, 373-383



Exosome Nanovesicles

- Nanosize, abundant, and stable
- Crossing barrier
- Home selection



Exosomes as Drug Delivery Carriers

- **Alternatives to traditional nanoparticles**
 - Natural carriers
 - No immunogenicity
 - Long circulating time
 - Cross physiological barrier
 - Active home-targeting
- **Borrowing self-secreted nanovesicles**
 - Cancer cells




Alvarez-Erviti et al (2011), Nature Biotechnology 29:341-5

Johnsen KB et al (2014), Biochimica et Biophysica Acta 1846:75-87

Hypothesis

- Cancer cell-derived exosomes could be used as a targeted carrier to deliver therapeutics into parental cancer cells




Experimental Design

- I. Isolate exosomes from cell media
 - A549 exosome: adenocarcinoma A549
 - bEND.3 exosome: brain endothelial bEND.3
- II. Assess home-selectivity of exosomes
- III. Test cell uptake of exosome-delivered fluorescent marker
- IV. Evaluate cytotoxicity of exosome-delivered anticancer drugs
- V. Identify knockdown efficacy of exosome-delivered siRNA on ICAM-1

I: Particle Sizing of Exosome

Exosome origin	Particle size
Brain endothelial bEND.3 cell	48.4 ± 3.3 nm
Lung adenocarcinoma A549 cell	25.5 ± 1.0 nm




II: Home Selectivity of Exosomes


- Exosome labeled by Exo-Green[®] kit
- **A549 cell** staining and imaging

bEND.3 exosome

Exosome Nucleus Overlay






X 4



X 40

A549 exosome

Exosome Nucleus Overlay



Home Selectivity of Exosomes


- Exosome labeled by Exo-Green[®] kit
- **bEND.3 cell** staining and imaging

bEND.3 exosome


Exosome

Nucleus

Overlay



X 4






X 40

A549 exosome

Exosome




Nucleus

Overlay



III: Cell Uptake of Rhodamine 123

- Loaded rhodamine 123 into exosomes
- A549 cancer cell
- Florescence microcopy
- C6 cell flow cytometry



I: Rhodamine 123

II: Rhodamine 123 loaded into A549 exosome



III: Rhodamine 123 loaded into bEND.3 exosome

* Results are significantly different, $p < 0.05$


IV: Exosome-Delivered Anticancer Drugs

- Doxorubicin (auto-red fluorescence)
- Cell uptake assessed by staining and imaging
- Doxorubicin, methotrexate, and paclitaxel formulated into different amounts of exosome
 - Formulation having more loaded drugs in the higher amount of exosome
- A549 cells treatment
- Cytotoxicity evaluation of exosome-delivered anticancer drugs


A549 Cell Uptake of Doxorubicin




Increased Cytotoxicity of Exosome-Delivered Doxorubicin




1: doxorubicin (1 μ M), 2. 200 μ g/ml exosome; 3. drug + 20 μ g/ml exosome; 4. drug + 40 μ g/ml exosome; 5. drug + 100 μ g/ml exosome; and 6. drug + 200 μ g/ml exosome, * Results are significantly different, $p < 0.05$).




Increased Cytotoxicity of Exosome-Delivered **Methotrexate**




1: methotrexate (50 μ M), 2. 200 μ g/ml exosome; 3. drug + 20 μ g/ml exosome; 4. drug + 40 μ g/ml exosome; 5. drug + 100 μ g/ml exosome; and 6. drug + 200 μ g/ml exosome, * Results are significantly different, $p < 0.05$).



Increased Cytotoxicity of Exosome-Delivered **Paclitaxel**




1. paclitaxel (50 μ M); 2. 200 μ g/ml exosome; 3. drug + 20 μ g/ml exosome; 4. drug + 40 μ g/ml exosome; 6. drug + 100 μ g/ml exosome; and 7. drug + 200 μ g/ml exosome, * Results are significantly different, $p < 0.05$).




V: Exosome-Delivered Small Interfering RNA (siRNA)

- Difficult to directly diffuse by incubation
- Form siRNA and transfection reagent (Lipofectamine[®]) complex
- Intercellular Adhesion Molecule 1 (ICAM-1)



ICAM-1 Knockdown by Selective SiRNA

ICAM-1 →



I. Buffer control

II. 60 µg A549 exosome


III. 60 nmole Scramble siRNA complex

IV. 60 nmole ICAM-1 siRNA complex

V. 60 nmole Scramble siRNA complex + 60 µg A549 exosome


VI. 60 nmole ICAM-1 siRNA complex + 60 µg A549 exosome

* Results are significantly different, $p < 0.05$



Inhibition From SiRNA Delivered by Autologous Cell-Derived Exosomes

ICAM-1 →



I. Buffer control

II. 60 nmole ICAM-1 siRNA


III. 60 nmole ICAM-1 siRNA complex

IV. 10 µl Transfection agent

V. 60 nmole ICAM-1 siRNA complex + 60 µg A549 exosome



VI. 60 nmole ICAM-1 siRNA complex + 60 µg bEND.3 exosome

* Results are significantly different, $p < 0.05$



Dose-dependent Effect of SiRNA Delivered by Exosomes

ICAM-1 →




I. Buffer control

II. 240 µg A549 exosome

III. 240 nmole ICAM-1 SiRNA

IV. 240 nmole ICAM-1SiRNA complex + 240 µg A549 exosome

* Results are significantly different, $p < 0.05$




Conclusions

- Nanosized A549 exosomes showed parental cancer cell homing-selectivity.
- High fluorescence intensity was observed in the cells treated with exosome-delivered rhodamine 123.
- Exosome-delivered anticancer drugs significantly enhanced cytotoxicity on the autologous cancer cells.
- Exosome-delivered siRNA presented dose-dependent knockdown efficacy in cancer cells. ICAM-1 inhibitions were based on exosome original sources and formulation methods.

Future Direction

- New types of exosome
- Safety profiles



Personalized treatment

Acknowledgments

➤ **Bai Lab**

Brittany Fogarty

Paige Martin

Ryan Hall

Joshua Allen

Alexandra Silber, Pharm.D.

Kaitlynn Pike, Pharm.D.

Raymond Bloom, Pharm.D.

➤ **Husson University**

Tianzhi Yang, Ph.D.

Roger Phipps, Ph.D.

➤ **University of Cincinnati**

Jiukuan Hao, Ph.D.

➤ **Mount Desert Island Biological Laboratory**

Voot Yin, Ph.D.

➤ **West Virginia University**

Paul Lockman, Ph.D.

➤ **Funding sources**



Questions

