Application Of Amorphous Solid Dispersion Technology For Improving The Solubility And In vitro dissolution properties Of *Withania somnifera* methanolic root powder extract



Kiran Dudhat*, R K School of Pharmacy, Rajkot, Gujarat, India

Abstract

Withania somnifera is a xerophytic plant whose methanolic extract of leaves and roots powder has exhibited effectiveness against various diseases. Like other natural products, its methanolic root powder extract also displayes poor dissolution because of high lipophilicity and poor aqueous solubility. The current reseach work was intended to formulate polymeric amorphous solid dispersion (PASD) system of methanolic extract of *Withania somnifera* root powder for improving its dissolution characteristics. The solvent evaporation method with three different polymers - HPMCAS-L, PVP K-30, and soluplus® were used to prepare the amorphous system. The complete transformation of the crystalline phytoconstituents to an amorphous/molecularly dispersed state was confirmed by DSC and PXRD studies, with all tested polymers at a 1:4 polymer ratio. The stability of phytoconstituents within prepared PASDs was confirmed by FTIR study, which revealed no changes in any of the distinctive peaks of pure extract. The prepared PASDs had significantly higher saturation solubility compared to pure extract. The formulated PASDs demonstrated substantial improvement in the dissolution properties of phytoconstituents compared to a pure extract containing capsules (f_2<50).

Bibliography

1. Gupta GL, Rana AC. Withania somnifera (Ashwagandha): a review. Pharmacognosy Reviews. 2007;1(1).

2. Kulkarni SK, Dhir A. Withania somnifera: an Indian ginseng. Progress in neuro-psychopharmacology and biological psychiatry. 2008;32(5):1093-105.

3. Devi PU, Sharada AC, Solomon FE. Antitumor and radiosensitizing effects of Withania somnifera (Ashwagandha) on a transplantable mouse tumor, Sarcoma-180. Indian journal of experimental biology. 1993;31(7):607-11.

4. Malik F, Singh J, Khajuria A, Suri KA, Satti NK, Singh S, et al. A standardized root extract of Withania somnifera and its major constituent withanolide-A elicit humoral and cell-mediated immune responses by up regulation of Th1-dominant polarization in BALB/c mice. Life Sciences. 2007;80(16):1525-38.
5. Owais M, Sharad KS, Shehbaz A, Saleemuddin M. Antibacterial efficacy of Withania somnifera (ashwagandha) an indigenous medicinal plant against experimental murine salmonellosis. Phytomedicine. 2005;12(3):229-35.
6. Chouhan N, Mittal V, Kaushik D, Khatkar A, Raina M. Self emulsifying drug delivery system (SEDDS) for phytoconstituents: a review. Current Drug Delivery. 2015;12(2):244-53.
7. Sayed N, Khurana A, Godugu C. Pharmaceutical perspective on the translational hurdles of phytoconstituents and strategies to overcome. Journal of Drug Delivery Science and Technology. 2019;53:101201.

Importance of Research and Development

Withania somnifera (native name - Ashwagandha) is known as "Indian ginseng" due to its broad therapeutical activities [1]. Various parts of the *Withania somnifera* shrub are used to treat a variety of diseases, such as adaptogenic, aphrodisiac, tonic, narcotic, diuretic, antihelminthic, astringent etc[2]. The broad range of therapeutical applicability of *Withania somnifera* is due to the presence of diverse secondary metabolites (steroidal alkaloids and lactones - a class of constituents together known as withanolides) present in different parts to plants (roots, stems, and leaves). The root, out of all the plant parts containing secondary metabolites, is considered to be the most therapeutically active as it contains a significant amount of phytoconstituents, especially withanolides. The methanolic extract of Withania somnifera root powder has displayed anti-tumor, immune-modulatory effects, anti-bacterial effects, anti-oxidant and anti-inflammatory activity [3-5]. However, like the majority of phytoconstituents, withanolides also exhibit extremely poor solubility and hence impaired dissolution, resulting in poor bioavailability after oral administration. After oral administration, adequate bioavailability is the primary concern for converting the phytoconstituents into successful market formulations, since these molecules are commonly used to treat several diseases, such as cancer, neurodegenerative diseases, infectious diseases, and chronic metabolic disorders, which require long-term treatment [6-7]. Another limitation to converting the alcoholic extract containing phytoconstituents into a successful oral market formulation is its nature of being sticky, hygroscopic, and adhesive, which is difficult to process at an industrial scale. Based on above-mentioned facts, the present research work was aimed to formulate and evaluated the ASD of *Withania somnifera* methanolic root powder extract to enhance its dissolution characteristics. The prepared amorhous solid dispersion (ASD) was also

evaluated for its ability to improve the *in-vitro* antioxidant activity of the methanolic root powder extract.

About University

RK University's philosophy is to educate through discovery. We shape thinkers & self-learners, making knowledge an exciting journey by opening new doors & steering into new horizons. Our students are a generation of tomorrow, a timeless tribe of change makers. When the fire burns within them, we rouse it, to scale greater heights. When ideas germinate, we support our students to become entrepreneurs and help to cultivate it, to bring about revolutions of Change. University offers verity of courses ranging from graduate level to post graduate levels such as BACHELOR OF TECHNOLOGY, BACHELOR OF SCIENCE, BACHELOR OF PHARMACY, MASTER OF TECHNOLOGY and MASTER OF PHARMACY etc.

Recent publications

1. Sourabhkumar Jain, Kiran Dudhat, M. M. Soniwala, DoE-Based Solid Self-microemulsifying Drug Delivery System (S-SMEDDS) Approach for Improving the Dissolution Properties of Raltegravir Potassium, Journal of Pharmaceutical Innovation.
2. M Bhalodiya, J Chavda, D Mori, N Patel, R Manek, K Dudhat, Formulation and evaluation of amorphous solid dispersion Boerhaavia diffusa methanolic root extract for improving dissolution properties, Journal of Drug Delivery Science and Technology, 66, 102740.
3. Mori Dhaval, Poonam Vaghela, Kajal Patel, Keshvi Sojitra, Mohini Patel, Sushma Patel, Kiran Dudhat, Sunny Shah, Ravi Manek, Ramesh Parmar, Lipid-based emulsion drug delivery systems—a comprehensive review, Drug Delivery and Translational Research, 12, pages1616–1639 (2022).
4. M Patel, D Mori, K Dudhat, S Shah, J Chavda, A Patel, Saturation Solubility and Dissolution Property Improvement of Albendazole by Salt Formation Approach, Journal of Pharmaceutical Innovation, (2021).
5. T Gusai, M Dhavalkumar, M Soniwala, K Dudhat, J Vasoya, J Chavda, Formulation and optimization of microsponge-loaded emulgel to improve the transdermal application of acyclovir—a DOE based approach, Drug Delivery and Translational Research 11 (5), 2009-2029.
6. A Dholakiya, K Dudhat, J Patel, D Mori, An integrated QbD based approach of SMEDDS and liquisolid compacts to simultaneously improve the solubility and processability of hydrochlorthiazide, Journal of Drug Delivery Science and Technology 61, 102162

References

1. Gupta GL, Rana AC. Withania somnifera (Ashwagandha): a review. Pharmacognosy Reviews. 2007;1(1).
2. Kulkarni SK, Dhir A. Withania somnifera: an Indian ginseng. Progress in neuro-psychopharmacology and biological psychiatry. 2008;32(5):1093-105.
3. Devi PU, Sharada AC, Solomon FE. Antitumor and radiosensitizing effects of Withania somnifera (Ashwagandha) on a transplantable mouse tumor, Sarcoma-180. Indian journal of experimental biology. 1993;31(7):607-11.
4. Malik F, Singh J, Khajuria A, Suri KA, Satti NK, Singh S, et al. A standardized root extract of Withania somnifera and its major constituent withanolide-A elicit humoral and cell-mediated immune responses by up regulation of Th1-dominant polarization in BALB/c mice. Life Sciences. 2007;80(16):1525-38.
5. Owais M, Sharad KS, Shehbaz A, Saleemuddin M. Antibacterial efficacy of Withania somnifera (ashwagandha) an indigenous medicinal plant against experimental murine salmonellosis. Phytomedicine. 2005;12(3):229-35.
6. Chouhan N, Mittal V, Kaushik D, Khatkar A, Raina M. Self emulsifying drug delivery system (SEDDS) for phytoconstituents: a review. Current Drug Delivery. 2015;12(2):244-53.
7. Sayed N, Khurana A, Godugu C. Pharmaceutical perspective on the translational hurdles of phytoconstituents and strategies to overcome. Journal of Drug Delivery Science and Technology. 2019;53:101201.
8. Curatolo, W., Nightingale, J.A., Herbig, S.M., 2009. Utility of hydroxypropylmethylcellulose acetate succinate (HPMCAS) for initiation and maintenance of drug supersaturation in the GI milieu. Pharmaceutical research 26(6), 1419-1431.
9. Dedroog, S., Pas, T., Vergauwen, B., Huygens, C., Van den Mooter, G., 2020. Solid-state analysis of amorphous solid dispersions: Why DSC and XRPD may not be regarded as stand-alone techniques. Journal of pharmaceutical and biomedical analysis 178, 112937.
10. Devi, P.U., Sharada, A.C., Solomon, F.E., 1993. Antitumor and radiosensitizing effects of Withania somnifera (Ashwagandha) on a transplantable mouse tumor, Sarcoma-180. Indian journal of experimental biology 31(7), 607-611.
11. Gangurde, A.B., Kundaikar, H.S., Javeer, S.D., Jaiswar, D.R., Degani, M.S., Amin, P.D., 2015. Enhanced solubility and dissolution of curcumin by a hydrophilic polymer solid dispersion and its insilico molecular modeling studies. Journal of Drug Delivery Science and Technology 29, 226-237.