Neuropsychological performance differences between two groups of probable-AD patients from different areas of Brazil

Analucy Aury Vieira de Oliveira[1],

ABSTRACT. During normal aging there are some cognitive and behavioral changes similar to those observed in a transitional state or mild cognitive impairment (MCI) and early onset dementia, making it challenging for health care professionals to reach an accurate and reliable diagnosis. Objective: The current study examined the performance of two different groups of patients diagnosed with probable Alzheimer's disease (AD) on a neuropsychological test battery. Methods: Twenty-two AD patients from Brasília-DF (AD1) and thirty-four AD patients from Palmas-TO, northern Brazil (AD2), were selected and a short neuropsychological battery administered. To verify the reliability of these previous diagnoses of AD, both groups of patients were compared with a group of healthy controls. Results: AD patients showed cognitive deficit but scores were lower for the AD2 group compared with the AD1 group considering the cut-off point. Notably, patients from the AD1 group were older ($p=0.004$) and had less formal education ($p<0.001$) than those from the AD2 group. Comparing different cognitive domains between AD groups, *post hoc* analysis showed that the AD1 group was characterized by deficits in episodic memory retrieval ($p<0.001$), semantic memory ($p<0.001$) and verbal fluency ($p<0.001$). In contrast, the AD2 group showed lower scores in attention ($p=0.007$), executive functioning ($p<0.001$) and working memory ($p<0.001$). Conclusion: This pattern suggests that the Palma group of patients had a neuropsychological profile that was inconsistent with AD. Although the results of this study have important clinical implications, the effects of age, education, and gender on cognitive performance should be explored further.

Key words: dementia, cognitive impairment, Alzheimer's disease, neuropsychological assessment, diagnosis.

ii. NO IMAGE

iii. REFERENCES

1. Brazilian Institute of Geography and Statistics. Demographic Census (IBGE - Instituto Brasileiro de Geografia e Estatística. Censo demográfico). Rio de Janeiro: IBGE, 2000.

2. Herrera E, Caramelli P, Silveira A, Nitrini R. Epidemiologic survey of dementia in a community-dwelling Brazilian population. Alzheimer Dis Assoc Disord 2002;16:103-108.

3. Nitrini, R. Diagnóstico de demência: avaliação clínica neuropsicológica e através da tomografia computadorizada por emissão de fóton único. Livre-Docência. Faculdade de Medicina da USP, 1993.

4. Nitrini R, Caramelli P, Bottino CM, Damasceno BP, Brucki SM, Anghinah R. Academia Brasileira de Neurologia. [Diagnosis of Alzheimer's disease in Brazil: diagnostic criteria and

auxiliary tests. Recommendations of the Scientific Department of Cognitive Neurology and Aging of the Brazilian Academy of Neurology]. Arq Neuropsiquiatr 2005;63: 713-719.

5. American Psychiatric Association. Diagnostic and statistical manual of mental disorders 4 ed. Washington DC: APA, 1994.

IV. Analucy Aury Vieira de Oliveira graduate at Psychology from Universidade Católica de Goiás (1992) and master's at M Ed Tefl from College St Mark St John University Of Exeter (2003). Has experience in Psychology, acting on the following subjects: psycholinguistics, acquisition, learning, adolescense, sexuality, memory, motivation, speaking, learning and depression and Cognitive Behavioral Therapy (CBT) suicide, mental disorders.

V. Analucy Aury Vieira de Oliveira

CONTACT NUMBER: 55 62 982330098

SESSION NAME; NEURODEGENATIVE DISEASES

ORAL COMUNICATION

PASSPORT NUMBER:

