Vegetable Consumption and Bladder Cancer Risk: a Pooled Analysis of Individual Patient Data of Six Cohort Studies

Maria E. Goossens[1], Maree Brinkman[2], Bas Bueno-de-Mesquita[3,4], Eric J. Grant[5], Piet van den Brandt[6], Elisabete Weiderpass[7,8,9,10], Emily White[11], Alicja Wolk[12], Frank Buntinx[1,13] , Maurice P. Zeegers[13,14,15]

[1] Department of General Practice, Katholieke Universiteit Leuven, ACHG-KU Leuven, Kapucijnenvoer 33, Blok J, bus 7001, 3000 Leuven, Belgium; [2] The Cancer Council Victoria, Melbourne, Australia; [3]Dt. for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM) Bilthoven, The Netherlands; [4]Dt. of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, London, United Kingdom; [5]Department of Epidemiology Radiation Effects Research Foundation, Hiroshima, Japan; [6]Department of Epidemiology, Schools for Oncology and Developmental Biology and Public Health and Primary Care, Maastricht University Medical Centre, Maastricht, The Netherlands; [7]Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden; [8]Department of Research, Cancer Registry of Norway, Institute of Population-Based Cancer Research, Oslo, Norway; [9]Genetic Epidemiology Group, Folkhälsan Research Center, Helsinki, Finland; [10] Department of Community Medicine, University of Tromsø, The Arctic University of Norway, Tromsø, Norway; [11]Fred Hutchinson Cancer Research Center, Seattle, WA, US; [12] Division of Nutritional Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; [13]CAPHRI School for Public Health and Primary Care, University of Maastricht, The Netherlands; [14]School of Cancer Sciences, University of Birmingham, Birmingham, UK; [15]NUTRIM School for Nutrition and Translational Research in Metabolism, University of Maastricht, Maastricht, The Netherlands.

Abstract

Background

Studies investigating the association between dietary factors and bladder cancer (BC) risk have reported inconsistent results, which may have been influenced by sex and smoking status. Inconsistencies in the published research may be overcome by a pooled analysis of individual patient data. Our aim was to investigate the association of vegetable intake, both total and individual vegetable consumption, on the risk of BC stratified by sex and smoking status.

Methods

The BLadder cancer, Epidemiology and Nutritional Determinants (BLEND) study collected individual patient data on vegetables from 4,143 bladder cancer cases and 655,505 non-cases from 6 cohort studies. The association of vegetables on the risk of bladder cancer was estimated by calculating odds ratios (ORs) and 95% confidence

intervals (CIs) using a one step multilevel mixed-effects logistic regression model adjusted for age, sex and smoking.

Results

We observed an inverse trend in BC risk with greater intake of total vegetables: each extra daily portion of vegetables was associated with a lower risk of BC by 6% (OR=0.94 (95%CI 0.92,0.96)). This risk reduction was stronger for women (OR=0.94 (95%CI 0.91,0.97)) than for men (OR=0.96 (95%CI 0.94,0.99)) and most pronounced for current smokers (OR=0.92 (95%CI 0.88,0.95)) in contrast with never smokers (OR=0.98 (95%CI 0.94,1.02)).

Conclusion

In this pooled analysis we found a consistent decrease in BC risk with each daily portion of total vegetables. We also observed differences in these trends by gender and smoking status.

Keywords

Bladder cancer, diet, vegetables, risk, pooled analysis, individual patient data

Recent publications

Goossens ME, Brinkman M, Zeegers MP, Buntinx F. Re: Is the Inverse Association Between Selenium and Bladder Cancer Due to Confounding by Smoking? American Journal of Epidemiology. 2015.

Goossens ME, Buntinx F, Zeegers MP. Aetiology, demographics and risk factors for bladder cancer. chapter in the Oxford Textbook of Urological Surgery, Oxford University Press (OUP) 2017 (in press).

Goossens ME, Buntinx F, Zeegers MP, Driessen JH, De Bruin ML, de Vries F. Influence of metformin intake on the risk of bladder cancer in type 2 diabetes patients. Br J Clin Pharmacol. 2015.

Goossens ME, Zeegers MP, Bazelier MT, De Bruin ML, Buntinx F, de Vries F. Risk of bladder cancer in patients with diabetes: a retrospective cohort study. BMJ open. 2015;5(6):e007470.

Goossens ME, Isa F, Brinkman M, Mak D, Reulen R, Wesselius A, et al. International pooled study on diet and bladder cancer: the bladder cancer, epidemiology and nutritional

determinants (BLEND) study: design and baseline characteristics. Arch Public Health. 2016;74:30.

Goossens ME, Zeegers MP, van Poppel H, Joniau S, Ackaert K, Ameye F, et al. Phase III randomised chemoprevention study with selenium on the recurrence of non-invasive urothelial carcinoma. The SELEnium and BLAdder cancer Trial. Eur J Cancer. 2016;69:9-18.

Bibliography

Mieke Goossens completed her master of medicine cum laude in 1990 at the university of Leuven and worked for more than 25 years as a general practitioner in Strombeek-Bever, Belgium.

In 2008 she started her PhD at KU Leuven on bladder cancer epidemiology. She initiated the Selenium and bladder cancer study (Seleblat study) to investigate the effect of seleniumyeast 200 mcg on the recurrence of bladder cancer in 14 Belgian hospitals and collected individual patient data from 18 case-control and six cohort studies from all over the world on diet and bladder cancer to investigate the association between diet and the risk of developing bladder cancer (The BLadder cancer, Epidemiology and Nutritional Determinants (BLEND) study).

She was a scientific researcher at the University of Birmingham, UK, and the University of Utrecht, The Netherlands.

She works as a scientific collaborator at the Belgium cancer center.