Pharmaceutical Regulatory Affairs: Open Access

10th International Conference on **Pharmacognosy and Medicinal Plants** November 21-22, 2022 | Las Vegas, USA

ISSN: 2167-7689

Volume 11 | ISSUE 3

Abstract (600 word limits)

Drug discovery a history of a pharmacognostic approach

Dr. Antonio Steardo MSc

Since the ancient Romans and ancient Greek used medical herbs, the pharmacognostic profiles of medicinal herbs guaranteed cures and remedies for minor daily health problems. Over the centuries, the tradition of medical herbs has allowed many simple therapies and complex diseases treatment. This gave birth to a part of modern pharmaceutical chemistry at the end of the 19th century. It started by the extraction of active ingredients from medicinal herbs. During the twentieth century, chemical modification led natural molecules to it characterises the discovery of new molecules. They were improving their pharmacodynamic and pharmacokinetic profile. Both the research and discovery of herbal medicine came from tradition. Even though their study ran into nefarious errors. During my pharmacologist career, I combined endogenous pharmacological modulation systems looking for new therapies. The discovery of many drugs derives from pharmacognostic research and the ethnobotanical tradition. There is no doubt that the chemistry of pharmacognostic products leads to the extraction of complex structures sometimes. Indeed, the most fascinating part is the historical testimonies on herbs studied over the centuries by those who practiced them. The oriental medicine, as well as the ancient Aesculapius, Galen, Paracelsus and Hippocrates, used herbal medicine. Its uses still leave some surprises to those who practice Drug Discovery in this sense.

Biography (200 word limit)

Since the ancient Romans and ancient Greek used medical herbs, the pharmacognostic profiles of medicinal herbs guaranteed cures and remedies for minor daily health problems. Over the centuries, the tradition of medical herbs has allowed many simple therapies and complex diseases treatment. This gave birth to a part of modern pharmaceutical chemistry at the end of the 19th century. It started by the extraction of active ingredients from medicinal herbs. During the twentieth century, chemical modification led natural molecules to it characterises the discovery of new molecules. They were improving their pharmacodynamic and pharmacokinetic profile. Both the research and discovery of herbal medicine came from tradition. Even though their study ran into nefarious errors. During my pharmacologist career, I combined endogenous pharmacological modulation systems looking for new therapies. The discovery of many drugs derives from pharmacognostic research and the ethnobotanical tradition. There is no doubt that the chemistry of pharmacognostic products leads to the extraction of complex structures sometimes. Indeed, the most fascinating part is the historical testimonies on herbs studied over the centuries by those who practiced them. The oriental medicine, as well as the ancient Aesculapius, Galen, Paracelsus and Hippocrates, used herbal medicine. Its uses still leave some surprises to those who practice Drug Discovery in this sense.

References (With Hyperlink)

1 “Cannabidiol in inflammatory bowel diseases: a brief overview.”
Esposito G, Filippis DD, Cirillo C, Iuvone T, Capoccia E, Scuderi C, Steardo A, Cuomo R, Steardo L.
Phytother Res. 2013 May; 27(5):633-6. doi: 10.1002/ptr.4781. Epub 2012 Jul 20. PMID: 22815234
Reviews.
2 “Cannabidiol in medicine: a review of its therapeutic potential in CNS disorders.”
Scuderi C, Filippis DD, Iuvone T, Blasio A, Steardo A, Esposito G.
Phytother Res. 2009 May; 23(5):597-602. doi: 10.1002/ptr.2625.
3 “Differential cannabinoid receptor expression during reactive gliosis: a possible implication for a no psychotropic neuroprotection.”
De Filippis D, Steardo A, D'Amico A, Scuderi C, Cipriano M, Esposito G, Iuvone T.
Scientific World Journal. 2009 Mar 31; 9:229-35. doi: 10.1100/tsw.2009.31.
PMID: 19347234 Reviews.
4 “Palmitoylethanolamide improves colon inflammation through an enteric glia/toll like receptor 4-dependent PPAR-α activation.”
Esposito G, Capoccia E, Turco F, Palumbo I, Lu J, Steardo A, Cuomo R, Sarnelli G, Steardo L.
Gut. 2014 Aug; 63(8):1300-12. doi: 10.1136/gutjnl-2013-305005. Epub 2013 Sep 30. PMID: 24082036 Reviews.
5 “Neurotensin as Modulator of Basal Ganglia- Thalamocortical Motor Circuit – Emerging Evidence for Neurotensin NTS1 Receptor as a Potential Target in Parkinson's Disease.”
By Luca Ferraro, Tiziana Antonelli, Sarah Beggiato, Maria Cristina Tomasini, Antonio Steardo, Kjell Fuxe and Sergio Tanganelli
Submitted: November 11th 2010 Reviewed: June 16th 2011 Published: February 8th 2012 DOI: 10.5772/21128
https://www.intechopen.com/chapters/27864

Organization / University Logo

